arXiv:1110.2935v1 [math.CO] 13 Oct 2011

Prime bound of a graph

Abderrahim Boussaïri[*] Pierre Ille[†][‡]

October 14, 2011

Abstract

Given a graph G, a subset M of $V(G)$ is a module of G if for each $v \in V(G) \smallsetminus M$, v is adjacent to all the elements of M or to none of them. For instance, $V(G)$, $\varnothing$ and $\{v\}$ ($v \in V(G)$) are modules of G called trivial. Given a graph G, $m(G)$ denotes the largest integer m such that there is a module M of G which is a clique or a stable set in G with $|M| = m$. A graph G is prime if $|V(G)| \geq 4$ and if all its modules are trivial. The prime bound of G is the smallest integer $p(G)$ such that there is a prime graph H with $V(H) \supseteq V(G)$, $H[V(G)] = G$ and $|V(H) \smallsetminus V(G)| = p(G)$. We establish the following. For every graph G such that $m(G) \geq 2$ and $\log_2(m(G))$ is not an integer, $p(G) = \lceil \log_2(m(G)) \rceil$. Then, we prove that for every graph G such that $m(G) = 2^k$ where $k \geq 1$, $p(G) = k$ or $k + 1$. Moreover $p(G) = k+1$ if and only if G or its complement admits 2^k isolated vertices. Lastly, we show that $p(G) = 1$ for every non-prime graph G such that $|V(G)| \geq 4$ and $m(G) = 1$.

Mathematics Subject Classifications (1991): 05C70, 05C69

Key words: Module; prime graph; prime extension; prime bound; modular clique number; modular stability number

1 Introduction

A *graph* $G = (V(G), E(G))$ is constituted by a *vertex set* $V(G)$ and an *edge set* $E(G) \subseteq \binom{V(G)}{2}$. Given a set S, $K_S = (S, \binom{S}{2})$ is the *complete* graph on S whereas $(S, \varnothing)$ is the *empty* graph. Let G be a graph. With each $W \subseteq V(G)$ associate the *subgraph* $G[W] = (W, \binom{W}{2} \cap E(G))$ of G induced by W. A graph H is an *extension* of G if $V(H) \supseteq V(G)$ and $H[V(G)] = G$. Given $p \geq 0$, a p-extension of G is an extension H of G such that $|V(H) \smallsetminus V(G)| = p$. The

---

[*] Faculté des Sciences Aïn Chock, Département de Mathématiques et Informatique, Km 8 route d'El Jadida, BP 5366 Maarif, Casablanca, Maroc; `aboussairi@hotmail.com`.

[†] Institut de Mathématiques de Luminy, CNRS – UMR 6206, 163 avenue de Luminy, Case 907, 13288 Marseille Cedex 09, France; `ille@iml.univ-mrs.fr`.

[‡] Centre de recherches mathématiques, Université de Montréal, Case postale 6128, Succursale Centre-ville, Montréal, Québec, Canada H3C 3J7.

complement of G is the graph $\overline{G} = (V(G), \binom{V(G)}{2} \setminus E(G))$. A subset W of $V(G)$ is a *clique* (respectively a *stable set*) in G if $G[W]$ is complete (respectively empty). The largest cardinality of a clique (respectively a stable set) in G is the *clique number* (respectively the *stability number*) of G, denoted by $\omega(G)$ (respectively $\alpha(G)$). Given $v \in V(G)$, the *neighbourhood* $N_G(v)$ of v in G is the family $\{w \in V(G) : \{v, w\} \in E(G)\}$. Its *degree* is $d_G(v) = |N_G(v)|$. We will consider N_G as a function from $V(G)$ in $2^{V(G)}$. A vertex v of G is *isolated* if $N_G(v) = \varnothing$. The family of isolated vertices of G is denoted by $\mathrm{Iso}(G)$.

We use the following notation. Let G be a graph. For $v \neq w \in V(G)$,

$$(v, w)_G = \begin{cases} 0 & \text{if } \{v, w\} \notin E(G), \\ 1 & \text{if } \{v, w\} \in E(G). \end{cases}$$

Given $W \subsetneq V(G)$, $v \in V(G) \setminus W$ and $i \in \{0, 1\}$, $(v, W)_G = i$ means $(v, w)_G = i$ for every $w \in W$. Given $W, W' \subsetneq V(G)$, with $W \cap W' = \varnothing$, and $i \in \{0, 1\}$, $(W, W')_G = i$ means $(w, W')_G = i$ for every $w \in W$. Given $W \subsetneq V(G)$ and $v \in V(G) \setminus W$, $v \sim_G W$ means that there is $i \in \{0, 1\}$ such that $(v, W)_G = i$. The negation is denoted by $v \not\sim_G W$.

Given a graph G, a subset M of $V(G)$ is a *module* of G if for each $v \in V(G) \setminus M$, we have $v \sim_G M$. For instance, $V(G)$, $\varnothing$ and $\{v\}$ ($v \in V(G)$) are modules of G called *trivial*. Clearly, if $|V(G)| \leq 2$, then all the modules of G are trivial. On the other hand, if $|V(G)| = 3$, then G admits a non-trivial module. A graph G is then said to be *prime* if $|V(G)| \geq 4$ and if all its modules are trivial. For instance, given $n \geq 4$, the *path* $(\{1, \ldots, n\}, \{\{p, q\} : |p - q| = 1\})$ is prime. Given a graph G, G and $\overline{G}$ share the same modules. Thus G is prime if and only if $\overline{G}$ is.

Let S be a set with $|S| \geq 2$. Given $p \geq 1$, consider a p-extension G of $\overline{K_S}$. If $|S| \geq 2^p$, then G is not prime. Indeed, for each $s \in S$, we have $N_G(s) \subseteq V(G) \setminus S$ because S is a stable set in G. So if $|S| > 2^p$, then $(N_G)_{\restriction S} : S \longrightarrow 2^{V(G) \setminus S}$ is not injective. Thus $\{s, t\}$ is a non-trivial module of G for $s \neq t \in S$ such that $N_G(s) = N_G(t)$. Furthermore, if $(N_G)_{\restriction S} : S \longrightarrow 2^{V(G) \setminus S}$ is injective and if $|S| = 2^p$, then there is $s \in S$ such that $N_G(s) = \varnothing$. Therefore $s \in \mathrm{Iso}(G)$ and $V(G) \setminus \{s\}$ is a non-trivial module of G. On the other hand, the following is well known and is easily verified (see Sumner [17, Theorem 2.45] and also Corollary 4 below). Given a set S with $|S| \geq 2$, $\overline{K_S}$ admits a prime $\lceil \log_2(|S| + 1) \rceil$-extension. This is extended to any graph by Brignall [2, Theorem 3.7] as follows.

Theorem 1. *A graph G, with $|V(G)| \geq 2$, admits a prime extension H*

such that $|V(H) \setminus V(G)| \leq \lceil \log_2(|V(G)| + 1) \rceil$.

Following Theorem 1, we introduce the notion of prime bound. Let G be a graph. The *prime bound* of G is the smallest integer $p(G)$ such that G admits a prime $p(G)$-extension. Obviously $p(G) = 0$ when G is prime.

A prime extension H of G is *minimal* [14, 9, 18, 1, 10] if for every $W \subsetneq V(H)$ such that $H[W]$ is prime, $H[W]$ does not admit an induced subgraph isomorphic to G. Given a graph G, a prime $p(G)$-extension of G is clearly minimal. By Theorem 1, $p(G) \leq \lceil \log_2(|V(G)|+1) \rceil$. Observe also that $p(G) = p(\overline{G})$ for every graph G. By considering the clique number and the stability number, Brignall [2, Conjecture 3.8] conjectured the following.

Conjecture 1. *For each graph G with $|V(G)| \geq 2$,*

$$p(G) \leq \lceil \log_2(\max(\omega(G), \alpha(G)) + 1) \rceil.$$

We answer the conjecture positively by refining the notions of clique number and of stability number as follows. Given a graph G, the *modular clique number* of G is the largest integer $\omega_M(G)$ such that there is a module M of G which is a clique in G with $|M| = \omega_M(G)$. The *modular stability number* of G is $\alpha_M(G) = \omega_M(\overline{G})$. Obviously $\omega_M(G) \leq \omega(G)$ and $\alpha_M(G) \leq \alpha(G)$. For convenience, set

$$m(G) = \max(\alpha_M(G), \omega_M(G)).$$

We establish

Theorem 2. *For every graph G such that $m(G) \geq 2$,*

$$\lceil \log_2(m(G)) \rceil \leq p(G) \leq \lceil \log_2(m(G) + 1) \rceil.$$

On the one hand, it follows that $p(G) = \lceil \log_2(m(G)) \rceil$ for a graph G such that $m(G) \geq 2$ and $\log_2(m(G))$ is not an integer. On the other, if $\log_2(m(G))$ is a positive integer, that is, $m(G) = 2^k$ where $k \geq 1$, then $p(G) = k$ or $k+1$. We prove

Theorem 3. *For every graph G such that $m(G) = 2^k$ where $k \geq 1$,*

$$p(G) = k + 1 \text{ if and only if } |\mathrm{Iso}(G)| = 2^k \text{ or } |\mathrm{Iso}(\overline{G})| = 2^k.$$

Lastly, we show that $p(G) = 1$ for every non-prime graph G such that $|V(G)| \geq 4$ and $m(G) = 1$ (see Proposition 7).

The case of directed graphs is quite different. Recall that a *tournament* $T = (V(T), A(T))$ is a directed graph such that $|\{(v,w),(w,v)\} \cap A(T)| = 1$ for any $v \neq w \in V(T)$. For instance, the *3-cycle* $C_3 = (\{1,2,3\}, \{(1,2),(2,3),(3,1)\})$ is a tournament. A tournament T is *transitive* provided that for any $u, v, w \in V(T)$, if $(u,v),(v,w) \in A(T)$, then $(u,w) \in A(T)$. Given a tournament T, a subset I of $V(T)$ is an *interval* [16, 11] (or a *clan* [5]) of T if for any $x, y \in I$ and $v \in V(T) \setminus I$, we have: $(x,v) \in A(T)$ if and only if $(y,v) \in A(T)$. Once again, $V(T)$, $\varnothing$ and $\{v\}$ ($v \in V(T)$) are intervals of T called trivial. For instance, all the intervals of C_3 are trivial. On the other hand, a transitive tournament with at least 3 vertices admits a non-trivial interval. A tournament with at least 3 vertices is *indecomposable* [16, 11] (or *simple* [6, 7, 13]) if all its intervals are trivial. The *indecomposable bound* of a tournament T is defined as the prime bound of a graph. It is still denoted by $p(T)$.

Theorem 4 ([6, 7, 13])**.** *For a tournament T with $|V(T)| \geq 3$,*

$$p(T) \leq 2.$$

Moreover $p(T) = 2$ if and only if T is transitive and $|V(T)|$ is odd.

2 Preliminaries

We begin with the well known properties of the modules of a graph (for example, see [5, Lemma 3.4, Theorem 3.2, Lemma 3.9]).

Proposition 1. *Let G be a graph.*

1. *Given $W \subseteq V(G)$, if M is a module of G, then $M \cap W$ is a module of $G[W]$.*
2. *Given a module M of G, if N is a module of $G[M]$, then N is a module of G.*
3. *If M and N are modules of G, then $M \cap N$ is a module of G.*
4. *If M and N are modules of G such that $M \cap N \neq \varnothing$, then $M \cup N$ is a module of G.*
5. *If M and N are modules of G such that $M \setminus N \neq \varnothing$, then $N \setminus M$ is a module of G.*
6. *If M and N are modules of G such that $M \cap N = \varnothing$, then there is $i \in \{0,1\}$ such that $(M,N)_G = i$.*

Given a graph G, a partition P of $V(G)$ is a *modular partition* of P if each element of P is a module of G. Let P be such a partition. Given $M \neq N \in P$, there is $i \in \{0,1\}$ such that $(M,N)_G = i$ by the last assertion of Proposition 1. This justifies the following definition. The *quotient* of G by P is the graph G/P defined on $V(G/P) = P$ by $(M,N)_{G/P} = (M,N)_G$ for $M \neq N \in P$. We use the following properties of the quotient (for example, see [5, Theorems 4.1–4.3, Lemma 4.1]).

Proposition 2. *Given a graph G, consider a modular partition P of G.*

1. *Given $W \subseteq V(G)$, if $|W \cap X| = 1$ for each $X \in P$, then $G[W]$ and G/P are isomorphic.*
2. *For any module M of G, $\{X \in P : M \cap X \neq \varnothing\}$ is a module of G/P.*
3. *For any module Q of G/P, the union $\cup Q$ of the elements of Q is a module of G.*

Given a graph G, with each non-empty module M of G, associate the modular partition $P_M = \{M\} \cup \{\{v\} : v \in V(G) \setminus M\}$. Given $m \in M$, the corresponding quotient G/P_M is isomorphic to $G[(V(G) \setminus M) \cup \{m\}]$ by the first assertion of Proposition 2. To associate a unique quotient with any graph and to characterize the corresponding quotient, the following strengthening of the notion of module is introduced. Given a graph G, a module M of G is said to be *strong* provided that for every module N of G, we have: if $M \cap N \neq \varnothing$, then $M \subseteq N$ or $N \subseteq M$. We recall the following well known properties of the strong modules of a graph (for example, see [5, Theorem 3.3]).

Proposition 3. *Let G be a graph.*

1. *$V(G)$, $\varnothing$ and $\{v\}$, $v \in V$, are strong modules of G.*
2. *For a strong module M of G and for $N \subseteq M$, N is a strong module of G if and only if N is a strong module of $G[M]$.*

With each graph G, we associate the family $\Pi(G)$ of the maximal strong modules of G under inclusion among the proper and non-empty strong modules of G. The modular decomposition theorem is stated as follows.

Theorem 5 (Gallai [8, 12])**.** *For a graph G with $|V(G)| \geq 2$, the family $\Pi(G)$ realizes a modular partition of G. Moreover, the corresponding quotient $G/\Pi(G)$ is complete, empty or prime.*

Given a graph G with $|V(G)| \geq 2$, we denote by $\mathbb{S}(G)$ the family of the non-empty strong modules of G. As a direct consequence of the definition of a strong module, we obtain that the family $\mathbb{S}(G)$ endowed with inclusion, denoted by $(\mathbb{S}(G), \subseteq)$, is a tree called the *modular decomposition tree* [3] of G. Given $M \in \mathbb{S}(G)$ with $|M| \geq 2$, it follows from Proposition 3 that $\Pi(G[M]) \subseteq \mathbb{S}(G)$. Furthermore, given $W \subseteq V(G)$ (respectively $W \subsetneq V(G)$), $(\{M \in \mathbb{S}(G) : M \supseteq W\}, \subseteq)$ (respectively $(\{M \in \mathbb{S}(G) : M \supsetneq W\}, \subseteq)$) is a total order. Its smallest element is denoted by $W\uparrow$ (respectively $W\Uparrow$). By Proposition 3, if $M \in \mathbb{S}(G) \setminus \{V(G)\}$, then $M \in \Pi(G[M\Uparrow])$.

Let G be a graph with $|V(G)| \geq 2$. Using Theorem 5, we label $\mathbb{S}(G) \setminus \{\{v\} : v \in V(G)\}$ by the function λ_G defined as follows. For each $M \in \mathbb{S}(G)$ with $|M| \geq 2$,

$$\lambda_G(M) = \begin{cases} \blacksquare & \text{if } G[M]/\Pi(G[M]) \text{ is complete,} \\ \square & \text{if } G[M]/\Pi(G[M]) \text{ is empty,} \\ \sqcup & \text{if } G[M]/\Pi(G[M]) \text{ is prime.} \end{cases}$$



Figure 1: $\mathbb{S}(G) \setminus \{\{v\} : v \in V(G)\} = \{C, S, V(G)\}$, $\lambda_G(C) = \blacksquare$, $\lambda_G(S) = \square$ and $\lambda_G(V(G)) = \sqcup$.

In Figure 1, we depict a graph G defined on $V(G) = \{a, b\} \cup \{c_1, c_2, c_3\} \cup \{s_1, \dots, s_5\}$ by

$$\begin{cases} C = \{c_1, c_2, c_3\} \text{ is a clique in } G, \\ S = \{s_1, \dots, s_5\} \text{ is a stable set in } G, \\ (a, S)_G = (a, b)_G = (b, C)_G = 1, \\ (a, C)_G = (C, S)_G = (b, S)_G = 0. \end{cases}$$

For $i \in \{1, 2, 3\}$ and $j \in \{1, \dots, 5\}$, $G[\{a, b, c_i, s_j\}]$ is a path and hence is prime. Thus the non-trivial modules of G are the subsets M of C or of S with $|M| \geq 2$. It follows that

$$\mathbb{S}(G) = \{\{v\} : v \in V(G)\} \cup \{C, S, V(G)\}$$

$$\text{and } \begin{cases} \lambda_G(C) = \blacksquare, \\ \lambda_G(S) = \square, \\ \lambda_G(V(G)) = \sqcup. \end{cases}$$

3 Modular clique number and modular stability number

Given a non-prime graph G, we are looking for a prime extension H of G. We have to break the non-trivial modules of G as modules of H. Precisely, given a non-trivial module M of G, there must exist $v \in V(H) \setminus V(G)$ such that $v \not\sim_H M$. Moreover, we have only to consider the minimal non-trivial modules of G under inclusion.

Given a graph G, denote by $\mathcal{M}(G)$ the family of modules M of G such that $|M| \geq 2$, and denote by $\mathcal{M}_{\min}(G)$ the family of the minimal elements of $\mathcal{M}(G)$ under inclusion.

Lemma 1. *Let G be a graph. For every $M \in \mathcal{M}(G)$, $M \in \mathcal{M}_{\min}(G)$ if and only if either $|M| = 2$ or $G[M]$ is prime.*

Proof. To begin, consider $M \in \mathcal{M}_{\min}(G)$. It follows from the second assertion of Proposition 1 that all the modules of $G[M]$ are trivial. Thus $G[M]$ is prime if $|M| \geq 4$. Assume that $|M| \leq 3$. Since a graph on 3 vertices admits a non-trivial module, we obtain $|M| = 2$.

Conversely, consider $M \in \mathcal{M}(G)$ such that $|M| = 2$ or $G[M]$ is prime. Clearly $M \in \mathcal{M}_{\min}(G)$ when $|M| = 2$. So assume that $G[M]$ is prime and consider $N \in \mathcal{M}(G)$ such that $N \subseteq M$. By the first assertion of Proposition 1, N is a module of $G[M]$. As $G[M]$ is prime, $N = M$ and hence $M \in \mathcal{M}_{\min}(G)$. $\Diamond$

Let G be a graph. Following Lemma 1, we denote by $\mathcal{P}(G)$ the family of modules M of G such that $G[M]$ is prime. In Figure 1, $\mathcal{P}(G) = \varnothing$.

Lemma 2. *Let G be a graph. For any $P \in \mathcal{P}(G)$ and $M \in \mathcal{M}(G)$, either $P \cap M = \varnothing$ or $P \subseteq M$.*

Proof. Assume that $P \cap M \neq \varnothing$. By the first assertion of Proposition 1, $P \cap M$ is a module of $G[P]$. Since $G[P]$ is prime, either $|P \cap M| = 1$ or $P \cap M = P$. Suppose for a contradiction that $|P \cap M| = 1$. As $|M| \geq 2$, $M \smallsetminus P \neq \varnothing$. By the second to last assertion of Proposition 1, $P \smallsetminus M$ is a module of G. By the first assertion, $P \smallsetminus M$ would be a non-trivial module of $G[P]$. It follows that $P \cap M = P$. ◇

Given a graph G, consider $u \neq v \in V(G)$ and $v \neq w \in V(G))$ such that there are $M_{\{u,v\}}, M_{\{v,w\}} \in \mathcal{M}_{\min}(G)$ with $u, v \in M_{\{u,v\}}$ and $v, w \in M_{\{v,w\}}$. First, assume that $M_{\{u,v\}} \in \mathcal{P}(G)$. By Lemma 2, $M_{\{u,v\}} \subseteq M_{\{v,w\}}$ and hence $M_{\{u,v\}} = M_{\{v,w\}}$ because $M_{\{v,w\}} \in \mathcal{M}_{\min}(G)$. Similarly $M_{\{u,v\}} = M_{\{v,w\}}$ when $M_{\{v,w\}} \in \mathcal{P}(G)$. Second, assume that $\left|M_{\{u,v\}}\right| = \left|M_{\{v,w\}}\right| = 2$, that is, $M_{\{u,v\}} = \{u, v\}$ and $M_{\{v,w\}} = \{v, w\}$. By interchanging G and $\overline{G}$, assume that $\{u, v\}$ is a stable set in G. We obtain $(u, \{v, w\})_G = 0$ and hence $(w, \{u, v\})_G = 0$. Thus $\{u, v, w\}$ is a stable set in G. Furthermore $\{u, v, w\}$ is a module of G by the fourth assertion of Proposition 1. Since $\{u, v, w\}$ is a stable set in G, $\{u, w\}$ is a module of $G[\{u, v, w\}]$. By the second assertion of Proposition 1, $\{u, w\}$ is a module of G. By Lemma 1, $\{u, w\} \in \mathcal{M}_{\min}(G)$. In both cases, there exists $M_{\{u,w\}} \in \mathcal{M}_{\min}(G)$ such that $u, w \in M_{\{u,w\}}$. Consequently, the binary relation $\approx_G$ defined on $V(G)$ by

$$u \approx_G v \text{ if } \begin{cases} u = v \\ \text{or} \\ u \neq v \text{ and there is } M_{\{u,v\}} \in \mathcal{M}_{\min}(G) \text{ with } u, v \in M_{\{u,v\}}, \end{cases}$$

for any $u, v \in V(G)$, is an equivalence relation. The family of the equivalence classes of $\approx_G$ is denoted by $\mathfrak{M}(G)$. In Figure 1, $\mathfrak{M}(G) = \{C, S, \{a\}, \{b\}\}$.

Lemma 3. *Let G be a graph. For every $C \in \mathfrak{M}(G)$ such that $|C| \geq 2$, one and only one of the following holds*

- *$C \in \mathcal{P}(G)$,*
- *C is a maximal module of G under inclusion among the modules of G which are cliques in G,*
- *C is a maximal module of G under inclusion among the modules of G which are stable sets in G.*

Proof. Consider $C \in \mathfrak{M}(G)$ such that $|C| \geq 2$. Clearly C satisfies at most one of the three assertions above. For any $u \neq v \in C$, there is $M_{\{u,v\}} \in \mathcal{M}_{\min}(G)$ such that $u, v \in M_{\{u,v\}}$. By Lemma 1, either $\left|M_{\{u,v\}}\right| = 2$ or $M_{\{u,v\}} \in \mathcal{P}(G)$.

First, assume that there are $u \neq v \in C$ such that $M_{\{u,v\}} \in \mathcal{P}(G)$. Let $w \in C \smallsetminus \{u, v\}$. By Lemma 2, $M_{\{u,v\}} \subseteq M_{\{u,w\}}$ and hence $M_{\{u,v\}} = M_{\{u,w\}}$ because $M_{\{u,w\}} \in \mathcal{M}_{\min}(G)$. Thus $C = M_{\{u,v\}} \in \mathcal{P}(G)$.

Second, assume that $M_{\{u,v\}} = \{u, v\}$ for any $u \neq v \in C$. Given $u \neq v \in C$, assume that $\{u, v\}$ is a stable set in G by interchanging G and $\overline{G}$. As observed above, $\{u, v, w\}$ is a module of G and a stable set in G for every $w \in C \setminus \{u, v\}$. Therefore C is a stable set in G and it follows from the fourth assertion of Proposition 1 that C is a module of G. Furthermore consider $u \in C$ and $x \in V(G) \setminus C$. Since $u \not\approx_G x$, $\{u, x\}$ is not a module of G and hence there is $y \in V(G) \setminus \{u, x\}$ such that $y \not\sim_G \{u, x\}$. If $y \notin C$, then $y \not\sim_G C \cup \{x\}$ and $C \cup \{x\}$ is no longer a module of G. If $y \in C$, then $(y, x)_G = 1$ because $(y, u)_G = 0$. Thus $C \cup \{x\}$ is no longer a stable set in G. Consequently C is a maximal module of G among the modules of G which are stable sets in G. ◇

Let G be a graph. Following Lemma 3, denote by $\mathcal{C}(G)$ the family of the maximal elements of $\mathcal{M}(G)$ under inclusion among the elements of $\mathcal{M}(G)$ which are cliques in G, and denote by $\mathcal{S}(G)$ the family of the maximal elements of $\mathcal{M}(G)$ under inclusion among the elements of $\mathcal{M}(G)$ which are stable sets in G. In Figure 1, $\mathcal{C}(G) = \{C\}$ and $\mathcal{S}(G) = \{S\}$. The next is a simple consequence of Lemma 3.

Corollary 1. *For a graph G, $\{C \in \mathfrak{M}(G) : |C| \geq 2\} = \mathcal{C}(G) \cup \mathcal{S}(G) \cup \mathcal{P}(G)$.*

Proof. By Lemma 3,

$$\{C \in \mathfrak{M}(G) : |C| \geq 2\} \subseteq \mathcal{C}(G) \cup \mathcal{S}(G) \cup \mathcal{P}(G).$$

For the opposite inclusion, consider $C \in \mathcal{C}(G) \cup \mathcal{S}(G) \cup \mathcal{P}(G)$. First, assume that $C \in \mathcal{C}(G) \cup \mathcal{S}(G)$. By interchanging G and $\overline{G}$, assume that G is a clique of G. For any $c \neq d \in C$, $\{c, d\}$ is a module of $G[C]$. By the second assertion of Proposition 1, $\{c, d\}$ is a module of G. Thus $\{c, d\} \in \mathcal{M}_{\min}(G)$ and hence $c \approx_G d$. Therefore there is $D \in \mathfrak{M}(G)$ such that $D \supseteq C$. By Lemma 3, $D \in \mathcal{C}(G) \cup \mathcal{S}(G) \cup \mathcal{P}(G)$. If $D \in \mathcal{P}(G)$, then $D = C$ because $D \in \mathcal{M}_{\min}(G)$ by Lemma 1. So assume that $D \in \mathcal{C}(G) \cup \mathcal{S}(G)$. As C is a clique in G, $D \in \mathcal{C}(G)$ and $C = D$ by the maximality of $C \in \mathcal{C}(G)$.

Second, assume that $C \in \mathcal{P}(G)$. By Lemma 1, $C \in \mathcal{M}_{\min}(G)$ and hence $c \approx_G d$ for any $c \neq d \in C$. So there is $D \in \mathfrak{M}(G)$ such that $D \supseteq C$. By Lemma 3, $D \in \mathcal{C}(G) \cup \mathcal{S}(G) \cup \mathcal{P}(G)$. As $G[C]$ is prime, C is not included in a clique or a stable set in G. Therefore $D \in \mathcal{P}(G)$. By Lemma 1, $D \in \mathcal{M}_{\min}(G)$ and hence $C = D$. ◇

Given a graph G, set $\mathcal{I}(G) = \mathfrak{M}(G) \setminus (\mathcal{C}(G) \cup \mathcal{S}(G) \cup \mathcal{P}(G))$ and $I(G) = \{v \in V(G) : \{v\} \in \mathcal{I}(G)\}$. In Figure 1, $I(G) = \{a, b\}$.

Remark 1. Given a graph G, consider $M \in \mathcal{M}(G)$. There exists $N \in \mathcal{M}_{\min}(G)$ such that $N \subseteq M$. By considering $p \neq q \in N$, we obtain that there exist $p \neq q \in M$ such that $p \approx_G q$. By Corollary 1, there is $C \in \mathcal{C}(G) \cup \mathcal{S}(G) \cup \mathcal{P}(G)$ such that $|C \cap M| \geq 2$.

Let G be a graph. If $\omega_M(G) \geq 2$, that is, $\mathcal{C}(G) \neq \varnothing$, then $\omega_M(G) = \max(\{|C| : C \in \mathcal{C}(G)\})$. Similarly $\alpha_M(G) = \max(\{|C| : C \in \mathcal{S}(G)\})$ if $\alpha_M(G) \geq 2$. Consequently, if $m(G) \geq 2$, that is, $\mathcal{C}(G) \cup \mathcal{S}(G) \neq \varnothing$, then

$$m(G) = \max(\{|C| : C \in \mathcal{C}(G) \cup \mathcal{S}(G)\}).$$

Given a graph G, Sabidussi [15] introduced the following equivalence relation Sab_G on $V(G)$. Given $u, v \in V(G)$, $u \,\mathrm{Sab}_G\, v$ if $N_G(u) = N_G(v)$. If $\alpha_M(G) \geq 2$, then $\mathcal{S}(G)$ is the family of the equivalence classes of Sab_G which are not singletons.

We complete the section with another equivalence relation induced by the modular decomposition tree. It is used by Giakoumakis and Olariu [10] to construct a minimal prime extension of a graph. Given a graph G, consider the equivalence relation $\leftrightarrow_G$ defined on $V(G)$ by: given $v, w \in V(G)$, $v \leftrightarrow_G w$ if $\{v\}\twoheaduparrow = \{w\}\twoheaduparrow$. The set of the equivalence classes of $\leftrightarrow_G$ is denoted by $\mathfrak{S}(G)$. In Figure 1, $\mathfrak{S}(G) = \{C, S, \{a, b\}\}$.

Precisely, to construct a minimal prime extension of a graph G, Giakoumakis and Olariu [10] use only the elements of $\mathfrak{S}(G) \cap \mathcal{M}(G)$. The remainder of the section is mainly devoted to the study of $\mathfrak{S}(G) \cap \mathcal{M}(G)$ (see Proposition 5).

Lemma 4. *Let G be a graph.*

1. *Let $M \in \mathbb{S}(G)$ with $|M| \geq 2$. If $\{N \in \Pi(G[M]) : |N| = 1\} \neq \varnothing$, then $\{m \in M : \{m\} \in \Pi(G[M])\} \in \mathfrak{S}(G)$.*
2. *Let $C \in \mathfrak{S}(G)$ with $|C| \geq 2$. For every $c \in C$, $C\uparrow = \{c\}\twoheaduparrow$ and $C = \{c \in C\uparrow : \{c\} \in \Pi(G[C\uparrow])\}$.*
3. *For each $M \in \mathbb{S}(G)$ with $|M| \geq 2$, there is $C \in \mathfrak{S}(G) \cap \mathbb{S}(G)$ such that $|C| \geq 2$ and $C \subseteq M$.*

Proof. The first assertion follows from the definition of $\leftrightarrow_G$. For the second, consider $C \in \mathfrak{S}(G)$ with $|C| \geq 2$. For $c, d \in C$, we have $\{c\}\twoheaduparrow = \{d\}\twoheaduparrow$. Given $c_0 \in C$, we obtain $C \subseteq \{c_0\}\twoheaduparrow$ and hence $C\uparrow \subseteq \{c_0\}\twoheaduparrow$. As $|C| \geq 2$, we have also $\{c_0\} \subsetneq C\uparrow$ and hence $\{c_0\}\twoheaduparrow \subseteq C\uparrow$. Thus $C\uparrow = \{c\}\twoheaduparrow$ for every $c \in C$. It follows from the definition of $\leftrightarrow_G$ that for each $v \in V(G)$, $v \in C$ if and only if $\{v\}\twoheaduparrow = C\uparrow$. Furthermore it follows from the second assertion of Proposition 3 that for each $d \in C\uparrow$, $\{d\}\twoheaduparrow = C\uparrow$ if and only if $\{d\} \in \Pi(G[C\uparrow])$. Therefore $C = \{c \in C\uparrow : \{c\} \in \Pi(G[C\uparrow])\}$.

For the third assertion, consider a strong module M of G with $|M| \geq 2$. Let N be a minimal strong module of G under inclusion among the strong modules M' of G such that $|M'| \geq 2$ and $M' \subseteq M$. By the minimality of N, we obtain $\Pi(G[N]) = \{\{n\} : n \in N\}$. By the first assertion, $N \in \mathfrak{S}(G)$. ◇

Proposition 4. *For a graph G,*

1. $\mathcal{P}(G) = \{C \in \mathfrak{S}(G) \cap \mathbb{S}(G) : \lambda_G(C) = \sqcup\}$,
2. $\mathcal{C}(G) = \{C \in \mathfrak{S}(G) : |C| \geq 2, \lambda_G(C\uparrow) = \blacksquare\}$,
3. $\mathcal{S}(G) = \{C \in \mathfrak{S}(G) : |C| \geq 2, \lambda_G(C\uparrow) = \square\}$.

Proof. For the first assertion, consider $P \in \mathcal{P}(G)$. By Lemma 2, $P \in \mathbb{S}(G)$. By the third assertion of Lemma 4, there is $C \in \mathfrak{S}(G) \cap \mathbb{S}(G)$ such that $|C| \geq 2$ and $C \subseteq P$. As $P \in \mathcal{M}_{\min}(G)$ by Lemma 1, $C = P$. Since $G[P]$ is prime, $\Pi(G[P]) = \{\{p\} : p \in P\}$. By the first assertion of Lemma 4, $P \in \mathfrak{S}(G)$.

Moreover it follows from the first assertion of Proposition 2 that $G[P]/\Pi(G[P])$ is prime. Thus $\lambda_G(P) = \sqcup$.

Conversely, consider $C \in \mathfrak{S}(G) \cap \mathbb{S}(G)$ such that $\lambda_G(C) = \sqcup$. Clearly $C{\uparrow}= C$ because $C \in \mathbb{S}(G)$. As $C \in \mathfrak{S}(G)$, it follows from the second assertion of Lemma 4 that $\Pi(G[C]) = \{\{c\} : c \in C\}$. Furthermore we have $G[C]/\Pi(G[C])$ is prime because $\lambda_G(C) = \sqcup$. Thus $G[C]$ is prime as well by the first assertion of Proposition 2. Therefore $C \in \mathcal{P}(G)$.

For the second assertion, consider $C \in \mathcal{C}(G)$. Denote by Q the family of $M \in \Pi(G[C{\uparrow}])$ such that $M \cap C \neq \varnothing$. For each $M \in Q$, $C \smallsetminus M \neq \varnothing$ because $M \subsetneq C{\uparrow}$. As M is a strong module of G by Proposition 3, we obtain $M \subsetneq C$. Thus $|Q| \geq 2$ and $C = \cup Q$. Given $M \in Q$, consider $N \in Q \smallsetminus \{M\}$. For $m \in M$ and $n \in N$, $\{m,n\}$ is a module of $G[C]$ because $G[C]$ is complete. By the second assertion of Proposition 1, $\{m,n\}$ is a module of G. Since M is a strong module of G such that $m \in M \cap \{m,n\}$ and $n \in \{m,n\} \smallsetminus M$, we get $M \subseteq \{m,n\}$ and hence $M = \{m\}$. Thus $\{c\} \in \Pi(G[C{\uparrow}])$ for each $c \in C$. Set $D = \{c \in C{\uparrow} \colon \{c\} \in \Pi(G[C{\uparrow}])\}$. We have $C \subseteq D$ and $D \in \mathfrak{S}(G)$ by the first assertion of Lemma 4. Moreover, by the first assertion of Proposition 1, C is a module of $G[C{\uparrow}]$. It follows from the second assertion of Proposition 2 that $\{\{c\} : c \in C\}$ is a module of $G[C{\uparrow}]/\Pi(G[C{\uparrow}])$. Since $\{\{c\} : c \in C\}$ is a clique in $G[C{\uparrow}]/\Pi(G[C{\uparrow}])$, we obtain $\lambda_G(C{\uparrow}) = \blacksquare$. As $\lambda_G(C{\uparrow}) = \blacksquare$ and as $D = \{c \in C{\uparrow} \colon \{c\} \in \Pi(G[C{\uparrow}])\}$, $\{\{d\} : d \in D\}$ is a module of $G[C{\uparrow}]/\Pi(G[C{\uparrow}])$ and a clique in $G[C{\uparrow}]/\Pi(G[C{\uparrow}])$. Consequently D is a clique in G and D is a module of $G[C{\uparrow}]$ by the last assertion of Proposition 2. By the second assertion of Proposition 1, D is a module of G. Since $C \in \mathcal{C}(G)$, $C = D$.

Conversely, consider $C \in \mathfrak{S}(G)$ such that $|C| \geq 2$ and $\lambda_G(C{\uparrow}) = \blacksquare$. By the second assertion of Lemma 4, $C = \{c \in C{\uparrow} \colon \{c\} \in \Pi(G[C{\uparrow}])\}$. Since $\lambda_G(C{\uparrow}) = \blacksquare$, C is a clique in G and, as above for D, it follows from Propositions 1 and 2 that C is a module of G. Thus there is $D \in \mathcal{C}(G)$ such that $C \subseteq D$. As already proved, $D \in \mathfrak{S}(G)$ and hence $C = D$.

The third assertion follows from the second by interchanging G and $\overline{G}$. ◇

It follows from Proposition 4 that

Corollary 2. *For a graph G,*

$$\begin{cases} \omega_{\mathcal{M}}(G) = \max(\{|C| : C \in \mathfrak{S}(G), |C| \geq 2, \lambda_G(C{\uparrow}) = \blacksquare\}) \text{ if } \omega_{\mathcal{M}}(G) \geq 2, \\ \alpha_{\mathcal{M}}(G) = \max(\{|C| : C \in \mathfrak{S}(G), |C| \geq 2, \lambda_G(C{\uparrow}) = \square\}) \text{ if } \alpha_{\mathcal{M}}(G) \geq 2. \end{cases}$$

The next is a simple consequence of Corollary 1 and Proposition 4.

Corollary 3. *Let G be a graph.*

1. $\{C \in \mathfrak{M}(G) : |C| \geq 2\} \subseteq \{C \in \mathfrak{S}(G) : |C| \geq 2\}$.
2. *Given $C \in \mathfrak{S}(G)$ with $|C| \geq 2$,*

$$C \notin \mathfrak{M}(G) \text{ if and only if } C \neq C{\uparrow} \text{ and } \lambda_G(C{\uparrow}) = \sqcup.$$

3. *Let $C \in \mathfrak{S}(G) \smallsetminus \mathfrak{M}(G)$ with $|C| \geq 2$. For each $c \in C$, $\{c\} \in \mathfrak{M}(G) \smallsetminus \mathfrak{S}(G)$.*
4. *$I(G) = \{v \in V(G) : \{v\} \in \mathfrak{S}(G)\} \cup (\bigcup_{C \in \mathfrak{S}(G) \smallsetminus \mathfrak{M}(G)} \{\{c\} : c \in C\})$.*

Remark 2. The second assertion of Corollary 3 easily provides graphs G such that $\mathfrak{S}(G) \smallsetminus \mathfrak{M}(G) \neq \varnothing$. For instance, $\{a, b\} \in \mathfrak{S}(G) \smallsetminus \mathfrak{M}(G)$ in Figure 1.

Consider a prime graph G_0. Given $\alpha \notin V(G_0)$ and $u \in V(G_0)$, consider a 1-extension G of G_0 to $V(G_0) \cup \{\alpha\}$ such that $\{u, \alpha\}$ is a module of G, that is, $\alpha \in X_G(u)$ where $X = V(G_0)$. Clearly $\{u, \alpha\}$ is the single non-trivial module of G. Consequently

$$\begin{cases} \mathfrak{M}(G) = \{\{u, \alpha\}\} \cup \{\{v\} : v \in V(G_0) \smallsetminus \{u\}\}, \\ \mathbb{S}(G) = \{\{v\} : v \in V(G)\} \cup \{\{u, \alpha\}, V(G)\}. \end{cases}$$

Since $\mathbb{S}(G) = \{\{v\} : v \in V(G)\} \cup \{\{u, \alpha\}, V(G)\}$, we get

$$\mathfrak{S}(G) = \{\{u, \alpha\}, V(G_0) \smallsetminus \{u\}\}.$$

Thus $V(G_0) \smallsetminus \{u\} \in \mathfrak{S}(G) \smallsetminus \mathfrak{M}(G)$.

The following summarizes our comparison between $\mathfrak{M}(G)$ and $\mathfrak{S}(G)$ for a graph G.

Proposition 5. *For a graph G,*

$$\begin{aligned} \mathfrak{S}(G) \cap \mathcal{M}(G) &= \{C \in \mathfrak{M}(G) : |C| \geq 2\} \\ &= \mathcal{C}(G) \cup \mathcal{S}(G) \cup \mathcal{P}(G). \end{aligned}$$

Proof. By Corollary 1, $\{C \in \mathfrak{M}(G) : |C| \geq 2\} = \mathcal{C}(G) \cup \mathcal{S}(G) \cup \mathcal{P}(G)$. Furthermore, it follows from Proposition 4 that $\mathcal{C}(G) \cup \mathcal{S}(G) \cup \mathcal{P}(G) \subseteq \mathfrak{S}(G) \cap \mathcal{M}(G)$. So consider $C \in \mathfrak{S}(G) \cap \mathcal{M}(G)$. If $\lambda_G(C\uparrow) = \blacksquare$ or $\square$, then $C \in \mathcal{C}(G) \cup \mathcal{S}(G)$ by the last two assertions of Proposition 4. Thus assume that $\lambda_G(C\uparrow) = \sqcup$, that is, $G[C\uparrow]/\Pi(G[C\uparrow])$ is prime. By the second assertion of Lemma 4, $C = \{c \in C\uparrow : \{c\} \in \Pi(G[C\uparrow])\}$. Since C is a module of G, C is a module of $G[C\uparrow]$ by the first assertion of Proposition 1. Therefore $\{\{c\} : c \in C\}$ is a module of $G[C\uparrow]/\Pi(G[C\uparrow])$ by the second assertion of Proposition 2. As $G[C\uparrow]/\Pi(G[C\uparrow])$ is prime, $\{\{c\} : c \in C\} = \Pi(G[C\uparrow])$. Consequently, $C = C\uparrow$ and hence $C \in \mathcal{P}(G)$ by the first assertion of Proposition 4. ◇

Given a non-primitive and connected graph G, Giakoumakis and Olariu [10, Theorem 3.9] construct a minimal prime extension of G by adding $|C|-1$ vertices for each $C \in \mathcal{C}(G) \cup \mathcal{S}(G)$ and one vertex for each element of $\mathcal{P}(G)$.

4 Some prime extensions

We use the next corollary to prove Theorem 2.

Lemma 5. *Let S and S' be disjoint sets such that $|S'| = \lceil \log_2(|S|+1) \rceil \geq 2$, that is, $1 \leq 2^{|S'|-1} \leq |S| < 2^{|S'|}$. Consider any graph G defined on $V(G) = S \cup S'$ such that S and S' are stable sets in G.*

1. *Assume that $2^{|S'|-1} < |S| < 2^{|S'|}$. Then, G is prime if and only if*
$$\text{the function } (N_G)_{\restriction S} \text{ is an injection from } S \text{ into } 2^{S'} \smallsetminus \{\varnothing\}; \tag{4.1}$$
2. *Assume that $2^{|S'|-1} = |S|$. Then, G is prime if and only if* (4.1) *holds and for any $s \in S$ and $s' \in S'$,*
$$\text{if } d_G(s) = d_G(s') = 1, \text{ then } (s, s')_G = 0.$$

Proof. First, if there is $s \in S$ such that $N_G(s) = \varnothing$, then $s \in \mathrm{Iso}(G)$ and hence $V(G) \smallsetminus \{s\}$ is a non-trivial module of G. Second, if there are $s \neq t \in S$ such that $N_G(s) = N_G(t)$, then $\{s, t\}$ is a non-trivial module of G. Third, if there are $s \in S$ and $s' \in S'$ such that $N_G(s) = \{s'\}$ and $N_G(s') = \{s\}$, then $\{s, s'\}$ is a non-trivial module of G.

Conversely, assume that (4.1) holds. Consider a module M of G such that $|M| \geq 2$. We have to show that $M = V(G)$. As $(N_G)_{\restriction S}$ is injective, $M \nsubseteq S$, that is, $M \cap S' \neq \varnothing$.

For a first contradiction, suppose that $M \subseteq S'$. Recall that for each $s \in S$, either $M \cap N_G(s) = \varnothing$ or $M \subseteq N_G(s)$. Given $m \in M$, consider the function $f : S \longrightarrow 2^{((S' \smallsetminus M) \cup \{m\})} \smallsetminus \{\varnothing\}$ defined by

$$f(s) = \begin{cases} N_G(s) \text{ if } M \cap N_G(s) = \varnothing, \\ N_G(s) \cup \{m\} \text{ if } M \subseteq N_G(s), \end{cases}$$

for every $s \in S$. Since $(N_G)_{\restriction S}$ is injective, f is also and we would obtain that $|S| < 2^{|S'|-1}$. It follows that $M \cap S \neq \varnothing$.

For a second contradiction, suppose that $S' \smallsetminus M \neq \varnothing$. We have $(S \cap M, S' \smallsetminus M)_G = (S' \cap M, S' \smallsetminus M)_G = 0$. If $S \subseteq M$, then $(S, S' \smallsetminus M)_G = 0$ so that $(N_G)_{\restriction S}$ would be an injection from S into $2^{S' \cap M} \smallsetminus \{\varnothing\}$ which contradicts $|S| \geq 2^{|S'|-1}$. Thus $S \smallsetminus M \neq \varnothing$. We obtain $(S \smallsetminus M, S' \cap M)_G = (S \smallsetminus M, S \cap M)_G = 0$. As $(S \cap M, S' \smallsetminus M)_G = 0$, $(N_G)_{\restriction (S \cap M)} : S \cap M \longrightarrow 2^{S' \cap M} \smallsetminus \{\varnothing\}$ and hence $|S \cap M| \leq 2^{|S' \cap M|} - 1$. Since $(S \smallsetminus M, S' \cap M)_G = 0$, $(N_G)_{\restriction (S \smallsetminus M)} : S \smallsetminus M \longrightarrow 2^{S' \smallsetminus M} \smallsetminus \{\varnothing\}$ and hence $|S \smallsetminus M| \leq 2^{|S' \smallsetminus M|} - 1$. Therefore $|S| \leq 2^{|S' \cap M|} + 2^{|S' \smallsetminus M|} - 2 \leq 2^{|S'|-1}$. As $|S| \geq 2^{|S'|-1}$, we obtain $|S| = 2^{|S'|-1}$ and $2^{|S' \cap M|} + 2^{|S' \smallsetminus M|} = 2 + 2^{|S'|-1}$ so that $\min(|S' \cap M|, |S' \smallsetminus M|) = 1$. For instance, assume that $|S' \cap M| = 1$. Since $|S \cap M| \leq 2^{|S' \cap M|} - 1$, $|S \cap M| = 1$. There exist $s \in S$ and $s' \in S'$ such that $M = \{s, s'\}$. By what precedes, $(s, S' \smallsetminus \{s'\})_G = (s', S \smallsetminus \{s\})_G = 0$. As $N_G(s) \neq \varnothing$, we would obtain $N_G(s) = \{s'\}$, $N_G(s') = \{s\}$ and $(s, s')_G = 1$. It follows that $S' \subseteq M$.

Lastly, suppose that $S \smallsetminus M \neq \varnothing$. For each $s \in S \smallsetminus M \neq \varnothing$, we would have $(s, S')_G = (s, S \cap M)_G = 0$ and hence $N_G(s) = \varnothing$. It follows that $S \subseteq M$ and $M = S \cup S'$. ◇

Corollary 4. *Let S and S' be disjoint sets such that $|S| \geq 3$ and $|S'| = \lceil \log_2(|S|+1) \rceil$. There exists a prime graph G defined on $V(G) = S \cup S'$ and satisfying*

- *S and S' are stable sets in G;*
- *$(N_G)_{\restriction S} : S \longrightarrow 2^{S'} \smallsetminus \{\varnothing\}$ is injective;*
- *there exists an injection $\varphi_{S'} : S' \longrightarrow S$ such that $N_G(\varphi_{S'}(s')) = S' \smallsetminus \{s'\}$ for each $s' \in S'$.*



Figure 2: Corollary 4 when $S = \{s_1, \ldots, s_5\}$, $S' = \{s'_1, s'_2, s'_3\}$ with $\varphi_{S'}(s'_1) = s_5$, $\varphi_{S'}(s'_2) = s_4$ and $\varphi_{S'}(s'_3) = s_3$.

Proof. (See Figure 2.) As $|S'| = \lceil \log_2(|S|+1) \rceil$, we have $2^{|S'|-1} \leq |S| < 2^{|S'|}$ and hence $|S'| \leq |S|$. Thus there exists a bijection $\psi_{S'}$ from S' onto $S'' \subseteq S$. Consider the injection $f_{S''} : S'' \longrightarrow 2^{S'} \smallsetminus \{\varnothing\}$ defined by $s'' \mapsto S' \smallsetminus \{(\psi_{S'})^{-1}(s'')\}$. Let f_S be any injection from S into $2^{S'} \smallsetminus \{\varnothing\}$ such that $(f_S)_{\restriction S''} = f_{S''}$. Lastly, consider the graph G defined on $V(G) = S \cup S'$ such that S and S' are stable sets in G and $(N_G)_{\restriction S} = f_S$. Before applying Lemma 5, assume that $|S| = 2^{|S'|-1}$. Since $|S| \geq 3$, $|S'| \geq 3$. For each $s' \in S'$, there are $t' \neq u' \in S' \smallsetminus \{s'\}$. We obtain $N_G(\psi_{S'}(t')) = S' \smallsetminus \{t'\}$ and $N_G(\psi_{S'}(u')) = S' \smallsetminus \{u'\}$. Therefore $s' \in N_G(\psi_{S'}(t')) \cap N_G(\psi_{S'}(u'))$. It follows that $d_G(s') \geq 2$ for every $s' \in S'$. By Lemma 5, G is prime. ◇

We use the following two results to prove Theorem 2 when $\mathcal{P}(G) \neq \varnothing$. Given a graph G, consider $X \subsetneq V(G)$ such that $G[X]$ is prime. We utilize the following subsets of $V(G) \smallsetminus X$ (for instance, see [4, Lemma 5.1])

- $\mathrm{Ext}_G(X)$ is the set of $v \in V(G) \smallsetminus X$ such that $G[X \cup \{v\}]$ is prime;
- $\langle X \rangle_G$ is the set of $v \in V(G) \smallsetminus X$ such that X is a module of $G[X \cup \{v\}]$;
- for $u \in X$, $X_G(u)$ is the set of $v \in V(G) \smallsetminus X$ such that $\{u, x\}$ is a module of $G[X \cup \{v\}]$.

The family $\{\mathrm{Ext}_G(X), \langle X\rangle_G\} \cup \{X_G(u) : u \in X\}$ is denoted by $p_{G[X]}$. The next lemma follows from Proposition 1.

Lemma 6. *Given a graph G, consider $X \subsetneq V(G)$ such that $G[X]$ is prime. The family $p_{G[X]}$ is a partition of $V(G) \setminus X$. Moreover, for each module M of G, one and only one of the following holds*

- *$X \subseteq M$ and $V(G) \setminus M \subseteq \langle X\rangle_G$;*
- *there is a unique $u \in X$ such that $M \cap X = \{u\}$ and $M \setminus \{u\} \subseteq X_G(u)$;*
- *$M \cap X = \varnothing$ and M is included in an element of $p_{G[X]}$. Moreover, for $v, w \in M$, the function $X \cup \{v\} \longrightarrow X \cup \{w\}$, defined by $v \mapsto w$ and $u \mapsto u$ for $u \in X$, is an isomorphism from $G[X \cup \{v\}]$ onto $G[X \cup \{w\}]$.*

Lemma 7. *Let G be a prime graph. For every $\alpha \notin V(G)$, there are*

$$2^{|V(G)|} - 2|V(G)| - 2$$

distinct prime 1-extensions of G to $V(G) \cup \{\alpha\}$.

Proof. Consider any graph H defined on $V(H) = V(G) \cup \{\alpha\}$ such that $H[V(G)] = G$ and

$$N_H(\alpha) \in 2^{V(G)} \setminus (\{\varnothing, V(G)\} \cup \{N_G(v) : v \in V(G)\} \cup \{N_G(v) \cup \{v\} : v \in V(G)\}).$$

We verify that H is prime. Set $X = V(G)$. We have $H[X] = G$ is prime. If $\alpha \in \langle X\rangle_H$, then $N_H(\alpha) = \varnothing$ or $V(G)$. Thus $\alpha \notin \langle X\rangle_H$. If there is $v \in V(G)$ such that $\alpha \in X_H(v)$, then $N_H(\alpha) = N_G(v)$ or $N_G(v) \cup \{v\}$. Therefore $\alpha \notin X_H(v)$ for every $v \in V(G)$. It follows from Lemma 6 that $\alpha \in \mathrm{Ext}_H(X)$, that is, H is prime. Consequently the number of prime 1-extensions of G to $V(G) \cup \{\alpha\}$ equals

$$|2^{V(G)} \setminus (\{\varnothing, V(G)\} \cup \{N_G(v) : v \in V(G)\} \cup \{N_G(v) \cup \{v\} : v \in V(G)\})|.$$

Clearly $\varnothing \notin \{N_G(v) \cup \{v\} : v \in V(G)\}$, $V(G) \notin \{N_G(v) : v \in V(G)\}$ and $\{N_G(v) : v \in V(G)\} \cap \{N_G(v) \cup \{v\} : v \in V(G)\} = \varnothing$. Moreover, if there is $v \in V(G)$ such that $N_G(v) = \varnothing$ or $V(G) \setminus \{v\}$, then $V(G) \setminus \{v\}$ would be a non-trivial module of G. If there are $v \neq w \in V(G)$ such that $N_G(v) = N_G(w)$ or $N_G(v) \cup \{v\} = N_G(w) \cup \{w\}$, then $\{v, w\}$ would be a non-trivial module of G. As G is prime, $V(G) \notin \{N_G(v) \cup \{v\} : v \in V(G)\}$, $\varnothing \notin \{N_G(v) : v \in V(G)\}$ and for $v \neq w \in V(G)$, we have $N_G(v) \neq N_G(w)$ and $N_G(v) \cup \{v\} \neq N_G(w) \cup \{w\}$. Therefore $|2^{V(G)} \setminus (\{\varnothing, V(G)\} \cup \{N_G(v) : v \in V(G)\} \cup \{N_G(v) \cup \{v\} : v \in V(G)\})| = 2^{|V(G)|} - 2|V(G)| - 2$. ◇



Figure 3: Theorem 2 for the graph depicted in Figure 1.

5 Proofs of Theorems 2 and 3

Proof of Theorem 2. Consider a graph G such that $|m(G)| \geq 2$. To begin, we show that $p(G) \geq \lceil \log_2(m(G)) \rceil$. By interchanging G and $\overline{G}$, assume that there exists $S \in \mathcal{S}(G)$ with $|S| = m(G)$. Given an integer $p < \log_2(m(G))$, consider any p-extension H of G. We must prove that H is not prime. We have $2^{|V(H)\setminus V(G)|} < |S|$ so that the function $S \longrightarrow 2^{V(H)\setminus V(G)}$, defined by $s \mapsto N_H(s) \cap (V(H) \setminus V(G))$, is not injective. Thus there are $s \neq t \in S$ such that $N_H(s) \cap (V(H) \setminus V(G)) = N_H(t) \cap (V(H) \setminus V(G))$. In other words, $v \sim_H \{s, t\}$ for every $v \in V(H) \setminus V(G)$. As S is a module of G, we have $v \sim_G \{s, t\}$, that is, $v \sim_H \{s, t\}$ for every $v \in V(G) \setminus S$. Since S is a stable set in G, $(v, s)_H = (v, s)_G = (v, t)_G = (v, t)_H$. Therefore $\{s, t\}$ is a module of H and H is not prime.

To prove that $p(G) \leq \lceil \log_2(m(G) + 1) \rceil$, we must construct a prime $\lceil \log_2(m(G) + 1) \rceil$-extension H of G. Let S' be a set such that $S' \cap V(G) = \varnothing$ and $|S'| = \lceil \log_2(m(G) + 1) \rceil$. Let $s'_1 \in S'$. Consider $S_0 \in \mathcal{C}(G) \cup \mathcal{S}(G)$ such that $|S_0| = m(G)$, that is, $2^{|S'|-1} \leq |S_0| < 2^{|S'|}$. By interchanging G and $\overline{G}$, we can assume that S_0 is a clique or a stable set in G. (In Figure 3, $m(G) = 5$, $S_0 = S = \{s_1, \dots, s_5\} \in \mathcal{S}(G)$ and $S' = \{s'_1, s'_2, s'_3\}$.)

We consider any graph H defined on $V(G) \cup S'$ and satisfying the following.

1. S' is a stable set in H.
2. The subgraph $H[S_0 \cup S']$ of H is defined as follows
   - Assume that $|S_0| = 2$. We require that the subgraph $H[S_0 \cup S']$ of H is a path on 4 vertices and S_0 is a clique in $H[S_0 \cup S']$. Thus

$$d_{H[S_0 \cup S']}(s') = 1 \text{ for each } s' \in S'. \tag{5.1}$$

- Assume that $|S_0| \geq 3$. By Corollary 4, we can consider for $H[S_0 \cup S']$ a prime graph defined on $S_0 \cup S'$ such that

  $$\begin{cases} S_0 \text{ is a stable set in } H[S_0 \cup S']; \\ (N_{H[S_0 \cup S']})_{\restriction S_0} : S_0 \longrightarrow 2^{S'} \setminus \{\varnothing\} \text{ is injective;} \\ \text{there exists an injection } \varphi_{S'} : S' \longrightarrow S_0 \text{ with} \\ \qquad\qquad N_{H[S_0 \cup S']}(\varphi_{S'}(\, s')) = S' \setminus \{s'\} \text{ for each } s' \in S'. \end{cases} \tag{5.2}$$

  (In Figure 3, the subgraph $H[S_0 \cup S']$ is also depicted in Figure 2.)

Set $X = S_0 \cup S'$. In both cases, $H[X]$ is prime.

3. Let $C \in \mathcal{C}(G) \setminus \{S_0\}$. We have $|C| < 2^{|S'|}$. We consider for $H[C \cup S']$ a graph such that C is a clique in $H[C \cup S']$ and

   $$\left.\begin{aligned} f_C : C &\longrightarrow 2^{S'} \\ c &\longmapsto N_{H[C \cup S']}(c) \cap S' \end{aligned}\right\} \text{ is an injection from } C \text{ into } 2^{S'} \setminus \{S'\}. \tag{5.3}$$

   (In Figure 3, we have $C = \{c_1, c_2, c_3\} \in \mathcal{C}(G)$ and f_C is defined by

   $$c_1 \mapsto \{s'_2\},\ c_2 \mapsto \{s'_3\},\ c_3 \mapsto \varnothing.)$$

4. Let $S \in \mathcal{S}(G) \setminus \{S_0\}$. We have $|S| < 2^{|S'|}$. We consider for $H[S \cup S']$ a graph such that S is a stable set in $H[S \cup S']$ and

   $$\left.\begin{aligned} f_S : S &\longrightarrow 2^{S'} \\ s &\longmapsto N_{H[S \cup S']}(s) \end{aligned}\right\} \text{ is an injection from } S \text{ into } 2^{S'} \setminus \{\varnothing\}. \tag{5.4}$$

5. Let $P \in \mathcal{P}(G)$. As $G[P]$ is prime, it follows from Lemma 7 that $G[P]$ admits a prime 1-extension. We consider for $H[P \cup S']$ a graph such that

   $$\begin{cases} H[P] = G[P] \text{ is prime,} \\ \mathrm{Ext}_{H[P \cup S']}(P) = S' \text{ by using Lemma 7.} \end{cases} \tag{5.5}$$

6. Let $v \in I(G)$. Since S_0 is a module of G such that $v \notin S_0$, there is $i \in \{0, 1\}$ such that $(v, S_0)_G = i$. We consider for $H[\{v\} \cup S']$ a graph such that

   $$(v, s'_1)_{H[\{v\} \cup S']} \neq i. \tag{5.6}$$

   (In Figure 3, we have $S_0 = S$ and $I(G) = \{a, b\}$ with

   $$\begin{cases} (a, S)_G = 1 \text{ and } (a, s'_1)_{H[\{a\} \cup S']} = 0, \\ (b, S)_G = 0 \text{ and } (b, s'_1)_{H[\{b\} \cup S']} = 1.) \end{cases}$$

In the construction above, we have $H[N] = G[N]$ for each $N \in \mathfrak{M}(G)$. Thus we can also assume that $H[V(G)] = G$.

We begin with the following observation. For every module M of H such that $|M| \geq 2$, we have $M \cap S' \neq \varnothing$. Otherwise suppose that $M \subseteq V(G)$. By the first assertion of Proposition 1, $M \in \mathcal{M}(G)$. By Remark 1, there is $C \in \mathcal{C}(G) \cup \mathcal{S}(G) \cup \mathcal{P}(G)$ such that $|C \cap M| \geq 2$. As M is a module of H, it follows from the first assertion of Proposition 1 that $C \cap M$ is a module of $H[C \cup S']$. Since $H[S_0 \cup S']$ is prime, $C \neq S_0$. If $C \in (\mathcal{C}(G) \cup \mathcal{S}(G)) \setminus \{S_0\}$, then $(f_C)_{\restriction(C \cap M)}$ would be constant which contradicts (5.3) and (5.4). Lastly, suppose that $C \in \mathcal{P}(G)$. As $C \cap M$ is a module of $H[C] = G[C]$ by the first assertion of Proposition 1, it follows from Lemma 2 that $C \cap M = C$. Thus C would be a module of $H[C \cup S']$ so that $\langle C \rangle_{H[C \cup S']} = S'$ which contradicts (5.5) and Lemma 6. Consequently, $M \cap S' \neq \varnothing$ for every module M of H such that $|M| \geq 2$.

Now, we prove that H is prime. Consider a module M of H such that $|M| \geq 2$. We have to show that $M = V(H)$. As observed above, $M \cap S' \neq \varnothing$. By Lemma 6, either there is $s' \in S'$ such that $M \cap X = \{s'\}$ or $X \subseteq M$.

For a contradiction, suppose that there is $s' \in S'$ such that $M \cap X = \{s'\}$. By the first assertion of Proposition 1, $M \setminus \{s'\}$ is a module of G. By the last assertion of Proposition 1, there is $i \in \{0,1\}$ such that $(M \setminus \{s'\}, S_0)_G = i$. Thus $(s', S_0)_{H[S_0 \cup S']} = i$. Since S' is a stable set in $H[S_0 \cup S']$, we have $d_{H[S_0 \cup S']}(s') = 0$ or $d_{H[S_0 \cup S']}(s') \geq 2$ so that (5.1) does not hold. Therefore $|S_0| \geq 3$. Let $t' \in S' \setminus \{s'\}$. By (5.2), $(\varphi_{S'}(s'), s')_{H[S_0 \cup S']} = 0$ and $(\varphi_{S'}(t'), s')_{H[S_0 \cup S']} = 1$ which contradicts $(s', S_0)_{H[S_0 \cup S']} = i$. It follows that $X \subseteq M$.

First, consider $C \in \mathcal{C}(G) \setminus \{S_0\}$. Suppose for a contradiction that $C \cap M = \varnothing$. Thus $C \cap S_0 = \varnothing$ and it follows from the last assertion of Proposition 1 that there is $i \in \{0,1\}$ such that $(C, S_0)_G = i$. As $C \cap M = \varnothing$, we obtain $(C, M)_H = i$. In particular $(C, S')_{H[C \cup S']} = i$ and f_C would be constant which contradicts (5.3). Therefore $C \cap M \neq \varnothing$. Suppose for a contradiction that $C \setminus M \neq \varnothing$ and consider $c \in C \setminus M$. Since C is a clique of G, $(c, C \cap M)_G = 1$. Hence $(c, M)_H = 1$ and in particular $(c, S')_{H[C \cup S']} = 1$ which contradicts $f_C(c) \neq S'$. It follows that $C \subseteq M$. Similarly $S \subseteq M$ for every $S \in \mathcal{S}(G) \setminus \{S_0\}$.

Second, consider $P \in \mathcal{P}(G)$. By the first assertion of Proposition 1, $M \cap V(G)$ is a module of G. As $M \cap V(G) \supseteq S_0$, it follows from Lemma 2 that either $(M \cap V(G)) \cap P = \varnothing$ or $P \subseteq M \cap V(G)$. Suppose for a contradiction that $(M \cap V(G)) \cap P = \varnothing$. By the last assertion of Proposition 1, there is $i \in \{0,1\}$ such that $(P, S_0)_G = i$. As $S_0 \subseteq M$ and $M \cap P = \varnothing$, we obtain $(P, M)_H = i$. Thus $(P, S')_H = i$ and hence $\langle P \rangle_{H[P \cup S']} = S'$ which contradicts (5.5) and Lemma 6. It follows that $P \subseteq M$.

Lastly, it follows from (5.6) that $I(G) \subseteq M$. Consequently, $M = V(H)$. $\diamond$

Corollary 5. *For every graph G such that $|m(G)| \geq 2$, if $\log_2(m(G))$ is not an integer, then $p(G) = \lceil \log_2(m(G)) \rceil$.*

Proof. It suffices to apply Theorem 2 after recalling that $\lceil \log_2(m(G)) \rceil = \lceil \log_2(m(G) + 1) \rceil$ if and only if $\log_2(m(G))$ is not an integer. $\diamond$

Before showing Theorem 3, we observe

Lemma 8. *Given a graph G, if* $\mathrm{Iso}(G) \neq \varnothing$ *or* $\mathrm{Iso}(\overline{G}) \neq \varnothing$*, then*

$$p(G) \geq \lceil \log_2(\max(|\mathrm{Iso}(G)|, |\mathrm{Iso}(\overline{G})|) + 1) \rceil.$$

Proof. Let G be a graph such that $\max(|\mathrm{Iso}(G)|, |\mathrm{Iso}(\overline{G})|) > 0$. By interchanging G and $\overline{G}$, assume that $\mathrm{Iso}(G) \neq \varnothing$. Given $p < \lceil \log_2(|\mathrm{Iso}(G)| + 1) \rceil$, consider any p-extension H of G. We have $2^{|V(H) \smallsetminus V(G)|} \leq |\mathrm{Iso}(G)|$ and we verify that H is not prime.

For each $u \in \mathrm{Iso}(G)$, we have $N_H(u) \subseteq V(H) \smallsetminus V(G)$. Thus $(N_H)_{\restriction \mathrm{Iso}(G)}$ is a function from $\mathrm{Iso}(G)$ in $2^{V(H) \smallsetminus V(G)}$. As previously observed, if $(N_H)_{\restriction \mathrm{Iso}(G)}$ is not injective, then $\{u, v\}$ is a non-trivial module of H for $u \neq v \in \mathrm{Iso}(G)$ such that $N_H(u) = N_H(v)$. So assume that $(N_H)_{\restriction \mathrm{Iso}(G)}$ is injective. As $2^{|V(H) \smallsetminus V(G)|} \leq |\mathrm{Iso}(G)|$, we obtain that $(N_H)_{\restriction \mathrm{Iso}(G)}$ is bijective. Thus there is $u \in \mathrm{Iso}(G)$ such that $N_H(u) = \varnothing$, that is, $u \in \mathrm{Iso}(H)$. Therefore H is not prime. It follows that $p(G) \geq \lceil \log_2(\max(|\mathrm{Iso}(G)|, |\mathrm{Iso}(\overline{G})|) + 1) \rceil$. ◇

We prove Theorem 3 when $m(G) = 2$.

Proposition 6. *For every graph G such that $m(G) = 2$,*

$$p(G) = 2 \text{ if and only if } |\mathrm{Iso}(G)| = 2 \text{ or } |\mathrm{Iso}(\overline{G})| = 2.$$

Proof. By Theorem 2, $p(G) = 1$ or 2. To begin, assume that $|\mathrm{Iso}(G)| = 2$ or $|\mathrm{Iso}(\overline{G})| = 2$. By Lemma 8, $p(G) \geq 2$ and hence $p(G) = 2$. Conversely, assume that $p(G) = 2$. Let $\alpha \notin V(G)$. As $m(G) = 2$, $|C| = 2$ for each $C \in \mathcal{C}(G) \cup \mathcal{S}(G)$. Let $C_0 \in \mathcal{C}(G) \cup \mathcal{S}(G)$. We consider any graph H defined on $V(G) \cup \{\alpha\}$ and satisfying the following.

1. For each $C \in \mathcal{C}(G) \cup \mathcal{S}(G)$, $\alpha \not\sim_H C$.
2. Let $P \in \mathcal{P}(G)$. We have $G[P]$ is prime. Using Lemma 7, we consider for $H[P \cup \{\alpha\}]$ a prime 1-extension of $G[P]$ to $P \cup \{\alpha\}$.
3. Let $v \in I(G)$. There is $i \in \{0, 1\}$ such that $(v, C_0)_G = i$. We require that $(v, \alpha)_H \neq i$.
4. $H[V(G)] = G$.

Since $p(G) = 2$, H admits a non-trivial module M. First, we verify that $\alpha \in M$. Otherwise M is a module of G. By Remark 1, there is $C \in \mathcal{C}(G) \cup \mathcal{S}(G) \cup \mathcal{P}(G)$ such that $|C \cap M| \geq 2$. Suppose that $C \in \mathcal{C}(G) \cup \mathcal{S}(G)$. As $|C| = 2$, $C \subseteq M$ which contradicts $\alpha \not\sim_H C$. Suppose that $C \in \mathcal{P}(G)$. By Lemma 2, $C \cap M = C$. Thus C would be a module of $H[C \cup \{\alpha\}]$ which contradicts the fact that $H[C \cup \{\alpha\}]$ is prime. It follows that $\alpha \in M$.

Second, we show that $P \subseteq M$ for each $P \in \mathcal{P}(G)$. Since $H[P \cup \{\alpha\}]$ is prime and since $M \cap (P \cup \{\alpha\})$ is a module of $H[P \cup \{\alpha\}]$ with $\alpha \in M \cap (P \cup \{\alpha\})$, we obtain either $(M \smallsetminus \{\alpha\}) \cap P = \varnothing$ or $P \subseteq M \smallsetminus \{\alpha\}$. In the first instance, there is $i \in \{0, 1\}$ such that $(M \smallsetminus \{\alpha\}, P)_G = i$. Therefore $(\alpha, P)_H = i$ which contradicts the fact that $H[P \cup \{\alpha\}]$ is prime. It follows that $P \subseteq M$.

Third, we proof that $C \cap M \neq \varnothing$ for each $C \in \mathcal{C}(G) \cup \mathcal{S}(G)$. Otherwise consider $C \in \mathcal{C}(G) \cup \mathcal{S}(G)$ such that $C \cap M = \varnothing$. There is $i \in \{0,1\}$ such that $(M, C)_G = i$. Thus $(\alpha, C)_H = i$ which contradicts $\alpha \not\sim_H C$.

In particular we have $C_0 \cap M \neq \varnothing$. Let $v \in I(G)$. Since $(v, C_0 \cap M)_G \neq (v, \alpha)_H$, we obtain $v \in M$.

To conclude, consider $v \in V(H) \setminus M$. By what precedes, there is $C \in \mathcal{C}(G) \cup \mathcal{S}(G)$ such that $v \in C$. By interchanging G and $\overline{G}$, assume that $C \in \mathcal{S}(G)$. Since $v \sim_H M$ and $(v, C \cap M)_G = 0$, we obtain $(v, M)_H = 0$. In particular $(v, I(G))_G = 0$ and $(v, P)_G = 0$ for every $P \in \mathcal{P}(G)$. Let $D \in (\mathcal{C}(G) \cup \mathcal{S}(G)) \setminus \{C\}$. As $D \cap M \neq \varnothing$, we have $(v, D \cap M)_G = 0$ and hence $(v, D)_G = 0$ because D is a module of G. It follows that $v \in \mathrm{Iso}(G)$. Therefore $(C, V(G) \setminus C)_G = 0$ because C is a module of G. Since C is a stable set in G, we obtain $C \subseteq \mathrm{Iso}(G)$. Clearly $\mathrm{Iso}(G)$ is a module of G and a stable set in G. Thus $|\mathrm{Iso}(G)| \leq m(G) = 2$. Consequently $C = \mathrm{Iso}(G)$. $\diamond$

We use the following notation in the proof of Theorem 3. Given a graph G such that $m(G) \geq 3$, set

$$\mathcal{C}_{\max}(G) = \{C \in \mathcal{C}(G) : |C| = m(G)\},$$
$$\mathcal{S}_{\max}(G) = \{S \in \mathcal{S}(G) : |S| = m(G)\}.$$

Proof of Theorem 3. Consider a graph G such that $m(G) = 2^k$ where $k \geq 1$. By Theorem 2, $p(G) = k$ or $k+1$. To begin, assume that $|\mathrm{Iso}(G)| = 2^k$ or $|\mathrm{Iso}(\overline{G})| = 2^k$. By Lemma 8, $p(G) \geq k+1$ and hence $p(G) = k+1$.

Conversely, assume that $p(G) = k+1$. If $k = 1$, then it suffices to apply Proposition 6. So assume that $k \geq 2$. With each $C \in \mathcal{C}_{\max}(G) \cup \mathcal{S}_{\max}(G)$ associate $w_C \in C$. Set $W = \{w_C : C \in \mathcal{C}_{\max}(G) \cup \mathcal{S}_{\max}(G)\}$ and $G' = G[V(G) \setminus W]$.

We prove that $m(G') = 2^k - 1$. Given $C \in \mathcal{C}_{\max}(G) \cup \mathcal{S}_{\max}(G)$, $C \setminus \{w_C\}$ is a module of G' and $C \setminus \{w_C\}$ is a clique or a stable set in G'. Thus $2^k - 1 = |C \setminus \{w_C\}| \leq m(G')$. Consider $C' \in \mathcal{C}_{\max}(G') \cup \mathcal{S}_{\max}(G')$. We show that C' is a module of G. We have to verify that for each $C \in \mathcal{C}_{\max}(G) \cup \mathcal{S}_{\max}(G)$, $w_C \sim_G C'$. First, asume that there is $c \in (C \setminus \{w_C\}) \setminus C'$. We have $c \sim_G C'$. Furthermore $\{c, w_C\}$ is a module of G. Thus $w_C \sim_G C'$. Second, assume that $C \setminus \{w_C\} \subseteq C'$. Clearly $w_C \sim_G C'$ when $C' \subseteq C \setminus \{w_C\}$. Otherwise assume that $C' \setminus (C \setminus \{w_C\}) \neq \varnothing$. By interchanging G' and $\overline{G'}$, assume that C' is a clique in G'. As $C \setminus \{w_C\} \subseteq C'$ and $|C \setminus \{w_C\}| \geq 2$, we obtain that C is a clique in G. Since $(C \setminus \{w_C\}, C' \setminus C)_G = 1$ and since C is a module of G, we have $(w_C, C' \setminus C)_G = 1$. Furthermore $(w_C, C \setminus \{w_C\})_G = 1$ because C is a clique in G. Therefore $(w_C, C')_G = 1$. Consequently C' is a module of G. As C' is a clique or a stable set in G, there is $C \in \mathcal{C}(G) \cup \mathcal{S}(G)$ such that $C \supseteq C'$. If $C \notin \mathcal{C}_{\max}(G) \cup \mathcal{S}_{\max}(G)$, then $|C'| \leq |C| < m(G)$. If $C \in \mathcal{C}_{\max}(G) \cup \mathcal{S}_{\max}(G)$, then $C' \subseteq C \setminus \{w_C\}$ and hence $|C'| < |C| = m(G)$. In both cases, we have $|C'| = m(G') < m(G)$. It follows that $m(G') = 2^k - 1$.

By Corollary 5, $p(G') = k$ and hence there exists a prime k-extension H' of G'. We extend H' to $V(H') \cup W$ as follows. Let $C \in \mathcal{C}_{\max}(G) \cup \mathcal{S}_{\max}(G)$. Consider the function $f_C : C \setminus \{w_C\} \longrightarrow 2^{V(H') \setminus V(G')}$ defined by $c \mapsto N_{H'}(c) \setminus$

$V(G')$ for $c \in C \setminus \{w_C\}$. Since H' is prime, f_C is injective. As $|C \setminus \{w_C\}| = 2^k - 1$ and $|2^{V(H') \setminus V(G')}| = 2^k$, there is a unique $N_C \subseteq V(H') \setminus V(G')$ such that $f_C(c) \neq N_C$ for every $c \in C \setminus \{w_C\}$. Let H be the extension of H to $V(H') \cup W$ such that $N_H(w_C) \cap (V(H') \setminus V(G')) = N_C$ for each $C \in \mathcal{C}_{\max}(G) \cup \mathcal{S}_{\max}(G)$.

As $p(G) = k+1$, H is not prime. So consider a non-trivial module M of H. Set $X = V(H')$. We have $H[X]$ is prime. Given $u \in X$, we verify that $X_H(u) = \varnothing$. Otherwise there is $C \in \mathcal{C}_{\max}(G) \cup \mathcal{S}_{\max}(G)$ such that $w_C \in X_H(u)$. If $u \in V(G')$, then $\{u, w_C\}$ is a module of G. Therefore there is $D \in \mathcal{C}(G) \cup \mathcal{S}(G)$ such that $\{u, w_C\} \subseteq D$. Necessarily $D = C$ and we would obtain $f_C(u) = N_C$. So suppose that $u \in V(H') \setminus V(G')$. There is $i \in \{0,1\}$ such that $(w_C, C \setminus \{w_C\})_G = i$. Thus $(u, C \setminus \{w_C\})_{H'} = i$. Since f_C is injective, the function $g_C : C \setminus \{w_C\} \longrightarrow 2^{((V(H') \setminus V(G')) \setminus \{u\})}$, defined by $g_C(c) = f_C(c) \setminus \{u\}$ for $c \in C \setminus \{w_C\}$, is injective as well. We would obtain $2^k - 1 \leq 2^{k-1}$ which does not hold when $k \geq 2$. It follows that $X_H(u) = \varnothing$ for each $u \in X$. By Lemma 6, either $M \cap X = \varnothing$ or $X \subseteq M$. In the first instance, $M \subseteq W$ and M is a module of G which contradicts Remark 1. Consequently $X \subseteq M$. As M is a non-trivial module of H, there exists $C \in \mathcal{C}_{\max}(G) \cup \mathcal{S}_{\max}(G)$ such that $w_C \notin M$. By interchanging G and $\overline{G}$, assume that C is a stable set in G. We have $(w_C, C \setminus \{w_C\})_G = 0$ so that $(w_C, M)_H = 0$ and $(w_C, V(G'))_G = 0$. Given $D \in (\mathcal{C}_{\max}(G) \cup \mathcal{S}_{\max}(G)) \setminus \{C\}$, we obtain $(w_C, D \setminus \{w_D\})_G = 0$. Since D is a module of G, $(w_C, w_D)_G = 0$. It follows that $w_C \in \mathrm{Iso}(G)$. As at the end of the proof of Proposition 6, we conclude by $C = \mathrm{Iso}(G)$. ◇

Lastly, we examine the graphs G such that $m(G) = 1$. For these, $\mathcal{C}(G) = \mathcal{S}(G) = \varnothing$. Thus either $|V(G)| \leq 1$ or $|V(G)| \geq 4$ and G is not prime.

Proposition 7. *For every non-prime graph G such that $|V(G)| \geq 4$ and $m(G) = 1$, we have $p(G) = 1$.*

Proof. Since $m(G) = 1$, we have $\mathcal{C}(G) = \mathcal{S}(G) = \varnothing$. By Corollary 1, $\mathfrak{M}(G) = \mathcal{P}(G) \cup \mathcal{I}(G)$. By considering $V(G) \in \mathcal{M}(G)$, it follows from Remark 1 that there is $P_0 \in \mathcal{P}(G)$.

Let $\alpha \notin V(G)$. We consider any graph H defined on $V(G) \cup \{\alpha\}$ and satisfying the following.

1. Let $P \in \mathcal{P}(G)$. We have $G[P]$ is prime. Using Lemma 7, we consider for $H[P \cup \{\alpha\}]$ a prime graph such that $H[P] = G[P]$.

   Set $X = P_0 \cup \{\alpha\}$. We have $H[X]$ is prime.

2. Let $v \in I(G)$. Since P_0 is a module of G such that $v \notin P_0$, there is $i \in \{0,1\}$ such that $(v, P_0)_G = i$. We consider for $H[\{v, \alpha\}]$ the graph such that $(v, \alpha)_{H[\{v,\alpha\}]} \neq i$.

In the construction above, we have $H[N] = G[N]$ for each $N \in \mathfrak{M}(G)$. Thus we can also assume that $H[V(G)] = G$. As in the proof of Theorem 2, we verify the following. For every module M of H such that $|M| \geq 2$, we have $\alpha \in M$.

Now, we prove that H is prime. Consider a module M of H such that $|M| \geq 2$. By what precedes, $\alpha \in M$ and it follows from Lemma 6 that either $M \cap X = \{\alpha\}$

or $X \subseteq M$. In the first instance, there is $i \in \{0,1\}$ such that $(P_0, M \setminus \{\alpha\})_G = 1$. Thus $(P_0, \alpha)_H = 1$ which contradicts the fact that $H[P_0 \cup \{\alpha\}]$ is prime. It follows that $X \subseteq M$. We conclude as in the proof of Theorem 2. Since $(v, P_0)_G \neq (v, \alpha)_H$ for every $v \in I(G)$, we have $I(G) \subseteq M$. Lastly, consider $P \in \mathcal{P}(G) \setminus \{P_0\}$. As $H[P \cup \{\alpha\}]$ is prime, it follows from Proposition 1 that either $(M \setminus \{\alpha\}) \cap P = \varnothing$ or $P \subseteq M \setminus \{\alpha\}$. In the first instance, there is $i \in \{0,1\}$ such that $(P, P_0)_G = i$. Therefore $(P, \alpha)_H = 1$ which contradicts the fact that $H[P \cup \{\alpha\}]$ is prime. Consequently $P \subseteq M$. ◇

References

[1] A. Brandstädt, C.T. Hoàng, J.-M. Vanherpe, On minimal prime extensions of a four-vertex graph in a prime graph, Discrete Math. 288 (2004), 9–17.

[2] R. Brignall, Simplicity in relational structures and its application to permutation classes, Ph.D. Thesis, University of St Andrews, 2007.

[3] R. McConnell, F. de Montgolfier, Linear-time modular decomposition of directed graphs, Discrete Appl. Math. 145 (2005), 198–209.

[4] A. Ehrenfeucht, G. Rozenberg, Primitivity is hereditary for 2-structures, fundamental study, Theoret. Comput. Sci. 3 (70) (1990), 343–358.

[5] A. Ehrenfeucht, T. Harju, G. Rozenberg, The Theory of 2-Structures, A Framework for Decomposition and Transformation of Graphs, World Scientific, Singapore, 1999.

[6] P. Erdős, E. Fried, A. Hajnal, E.C. Milner, Some remarks on simple tournaments, Algebra Universalis 2 (1972), 238–245.

[7] P. Erdős, A. Hajnal, E.C. Milner, Simple one point extensions of tournaments, Mathematika 19 (1972), 57–62.

[8] T. Gallai, Transitiv orientierbare Graphen, Acta Math. Acad. Sci. Hungar. 18 (1967), 25–66.

[9] V. Giakoumakis, On the closure of graphs under substitution, Discrete Math. 177 (1997), 83–97.

[10] V. Giakoumakis, S. Olariu, All minimal prime extensions of hereditary classes of graphs, Theoret. Comput. Sci. 370 (2007), 74–93.

[11] P. Ille, Indecomposable graphs, Discrete Math. 173 (1997), 71–78.

[12] F. Maffray, M. Preissmann, "A translation of Tibor Gallai's paper: Transitiv orientierbare Graphen," Perfect Graphs J.L. Ramirez-Alfonsin and B.A. Reed, (Editors), Wiley, New York (2001), pp. 25–66.

[13] J.W. Moon, Embedding tournaments in simple tournaments, Discrete Math. 2 (1972), 389–395.

[14] S. Olariu, On the closure of triangle-free graphs under substitution, Inform. Process. Lett. 34 (1990), 97–101.

[15] G. Sabidussi, The composition of graphs, Duke Mathematical Journal 26 (1959), 693–696.

[16] J.H. Schmerl, W.T. Trotter, Critically indecomposable partially ordered sets, graphs, tournaments and other binary relational structures, Discrete Math. 113 (1993), 191–205.

[17] D.P. Sumner, Indecomposable graphs, Ph.D. Thesis, University of Massachusetts, 1971.

[18] I. Zverovich, Extension of hereditary classes with substitutions, Discrete Appl. Math. 128 (2003), 487–509.